

CORRELATION BETWEEN RADON DISTRIBUTION AND PREVALENCE OF LUNG CANCER IN WEST GEORGIA

1George I. MELIKADZE, 2Avtandil G. AMIRANASHVILI,
3Kakha G. GVINIANIDZE, 3David G. TSERETELI,
1Mariam Sh.TODADZE.

1Seismic Monitoring Center, 77, Nuthubidze Str., 0177, Tbilisi, Georgia,
melikadze@gmail.com

2Mikheil Nodia Institute of Geophysics, 1, M. Alexidze Str., 0193, Tbilisi, Georgia,
avto_amiranashvili@hotmail.com

3Institute of Public Health, 21, Kavtaradze Str, 0186, Tbilisi, Georgia
gvinianidze@hotmail.com

Abstract

Quantitative assessment of radon distribution in several regions of West Georgia has been carried out. According to field data in more than 100 water samples there is high content of radon. This research once again confirms correlation between Rn exhalation and prevalence of lung cancer.

Keywords: Rn mapping, lung cancer, indoor radon.

1. Introduction

According STCU project N 3992 “Assessment of radon-hazard potential, residential exposure, lung cancer and COPD in West Georgia” during 2007-2008 authors carried our field work in order to quantify the radon distribution, ascertain geological factors influencing indoor radon concentrations and find relation between radon content and health and in various geographical areas of West Georgia.

2. Method

When undertaking the gas survey of Rn the particular attention was paid to the multiple active zones of faults and areas of elevated geo-chemical background of uranium and quicksilver. Mobile group conducted the researches by Radon measurement PPA-01M-03 device. Rn content was measured in any type of water source (boreholes, wells and springs) and in the soil aeration zone in several regions ((Samtredia, Kutaisi, Vani, Tskaltubo, Khoni and Bagdadi). All observation sites were fixed by GPS measurements.

For ²²²Rn gas measurements in the dwellings the alpha track detectors (ATDs) were used to provide integrated (mean) radon concentration. Alpha track detectors were installed for the period of 12 months in the apartments and basements.

To reveal possible association between radon distribution and prevalence of lung cancer in West Georgia, we have performed an ecologic epidemiological study. Data on prevalence of lung cancer in various regions were collected through the primary health care net (primary health care net is represented by policlinics and out-patients' clinics), National Cancer Center and department of statistic of National Center of Disease Control and Public Health.

The key method for fulfillment the project is Radon mapping based on application of geochemical methods. Connection of anomalies to geological and hydro-geological structures, also medical data is analyzed using GIS technology.

3. Data analysis

Results of analyses on radon concentration were marked on topographic and geological maps. After that the field data were digitized and transferred into GIS-system. On the basis of these data the map of Rn content in water and soil were compiled using GIS technique (Fig.1). Areas of anomalously high Rn exhalation both in water and in soil were revealed in Tskhaltubo, Kuttaisi, Vani, Bagdadi, Chokhatauri and Ozurgeti regions. In order to understand the nature of these anomalies it is necessary to analyze all factors that influence intensity of Rn degassing and lead to high risk of its accumulation. These factors are geological structures of the region, presence of tectonic faults, presence of radioactive elements in the rocks, hydrogeological and geomorphological structures of the region, soil characteristics etc (Fig.2).

The Northern and Central parts of the tested area belongs from the geological point of view to the Kutaisi sub-zone of the Georgian plate and its Southern part – to the Adjara-Trialeti folded system. The North-East part of the first zone (villages Rioni and Gurna) is composed mostly by Jurassic volcanic rocks, which contain fissure groundwater of low mineralization (hydrocarbonate and calcium type). From the morphological point of view here we have glacial-erosion terrain that is characteristic for this mountainous area. In this area to the north of Kutaisi we found a band of elevated radon content in the soil (22-26 KBq/m³), which should be related to the presence of dikes of the crystalline rocks and systems of faults, developed on this territory. At the same time the content of radon in the water is low, which can be explained by influence of near surface groundwater circulation in this zone.



Fig. 1 Radon distribution in soil



Fig. 2 Geological map

To the South on the surface there are Lower Cretaceous rocks, which contain fissure and fissure-karstic type of pressurized ground water (regions of Tskaltubo and Kutaisi); the characteristic example is the low-radioactive thermal waters of Tskaltubo resort. Here the springs have large debit (200-220 l/s). The recharge of the aquifer takes place in the northern elevated areas; then the aquifer plunge under the Quarternary layers and its

discharge takes place at the contact area of Georgian plate and Adjara-Trialeti folded system, where a lot of transversal faults are found. This is also confirmed by the existence in this zone of low-radioactive thermal waters at the resorts Sulori, Amaghleba and Vani.

In the south and south-west part of test area, i.e. in the Adjara-Trialeti folded system (regions of Vani, Chokhatauri and Ozurgeti), in the volcanic and sediment rocks of Middle Eocen we observe karstic-fissure and fissure pressurized groundwaters of low radioactivity. The terrain here is of erosion-peneplain type.

Similar to Tskaltubo region here also are observed high values of Rn content in the soil (22-58 KBq/m³); this can be explained by high gas permeability of rocks and geomorphology of the area. As to the Rn content in water, it is a bit less (16-22 Bq/l) than in Tskaltubo region and cover much less area due to the fact that here mostly the shallow groundwater's are observed; these waters are characterized by shallow circulation system and they are not discharged on the surface (situation is alike to that in the North, where groundwater is in Volcanic rocks of Jurassic age). From the studied regions highest incidence rate (per 100.000 populations) of lung cancer is in several region (Tab. 1).

In regions with high Rn distribution on the whole is highest incidence rate of lung cancer (Figure 3).

Tab. 1 Distribution of parameters

Region	Incidence rate of lung cancer	Rn in Soil	ATDs
Thkaltubo	20,4	39.53	219.7
Bagdati	13,9	30.68	
Vani	14,7	34.78	112
Zestafoni	19,9	35.82	
Terdjola	15,5	31.46	
Samtredia	26,6	23.07	3.5
Khoni	38,5	35.81	
Zugdidi	23,2	19.54	
Abasha	32,0	15.81	
Martvili	26,9	47.72	
Senaki	25,0	13.85	
Lanchkhuti	49,0	76	
Ozurgeti	29,7	61	
Chokhatauri	57,0	90	




Fig. 3 Variation of parameters

In 2007 during field work in selected sites the alpha track detectors (ATDs) were installed; in 2008 part of them were dismantled and the tracks were investigated. The obtained results confirm the above conclusions. Increased long-time accumulation of Rn was observed in Tskhaltubo and Vani regions. Relatively less accumulation was in Samtredia region. In future the analysis of ATDs, installed in other regions will be continued.

4. Conclusions

Peculiarities of distribution of Rn on the territory of West Georgia was studied and anomalous areas were outlined. The elevated exhalation of Rn is the result of draining of Lower Cretaceous and Middle Eocen aquifers by rising springs and boreholes. This research once again confirms correlation between Rn exhalation and prevalence of lung cancer.

References

- [1] Amiranashvili A., Jishkariani D., Nodia A., Tatashidze Z. – Aeroionization condition and natural air radioactivity in Tskaltubo Cave, Int. J. of Immunorehabilitation, No1, suppl., 1994 a, 31-31
- [2] Auvinen A, Makelainen I, Hakama M, Castren O, Pukkala E, Reisbacka H et al. Indoor radon exposure and risk of lung cancer: a nested case-control study in Finland. J Natl Cancer Inst 1996;88:966-72 [Erratum: J Natl Cancer Inst 1998;90:401-2].
- [3] Field, RW. A review of residential radon case-control epidemiologic studies performed in the United States. Rev Environ Health 2001;16:151-67
- [4] Gil Hoon, Jai-Kilee – Construction of an environmental radon monitoring system using CR-39 nuclear track detectors, Nuclear Engineering and Technology, V. 37, N 4, 2005, 395-400
- [5] Health Effects of Exposure to Radon: Biological Effects of Ionizing Radiation (BEIR), 1999
- [6] Mellander H., Enflo A. - The Alpha Track Method Used in the Swedish Radon Epidemiological Study, Proc. of 5th Inter. Symposium on the Natural Radiation Environment, 1991, 22-28
- [7] Jonsson G. - Indoor 222Rn measurements in Sweden with the solid-state nuclear track detector technique, Health Phys., Vol. 54, No. 3, 1998, 271-281.
- [8] Krewski D, Lubin JH, Zielinski JM, Alavanja M, Catalan VS, Field RW, Klotz JB, Letourneau EG, Lynch CF, Lyon JJ, Sandler DP, Schoenberg JB, Steck DJ, Stolwijk JA, Weinberg C, Wilcox HB (2006) A Combined Analysis of North American Case-Control Studies of Residential Radon and Lung Cancer, Journal of Toxicology and Environmental Health, 69 (7-8), 533-597.

- [9] Kreienbrock L, Kreuzer M, Gerken MM, Dingerkus G, Wellmann J, Keller G et al. Case-control study on lung cancer and residential radon in Western Germany. *Am J Epidemiol* 2001;153:42-52.
- [10] Melikadze G.L., Adamchuk U.V., Buachidze G.I - "Radon correlation coefficient gravitation, displaying the seismic situation", 10 years after the Spitak earthquake Conference; materials, Yerevan 1998
- [11] Neuberger JS, Mahnken JD, Mayo MS, Field R. W. Risk Factors for Lung Cancer in Iowa Women: Implications for Prevention, *Cancer Detection and Prevention*, 2006; 30, 158-167
- [12] REMR – Radiation-Hygienic estimation and epidemiologic analysis of population irradiation levels in the Trans Caucasus, Radiation ecology of the mountainous region, (REMR), part I, Georgian Academy of Ecological Science, Tbilisi, 2006d, 219-235, (in Russian)